

DETECTION OF HARD X-RAY EMISSION FROM THE GALACTIC NUCLEAR REGION WITH *INTEGRAL* *

G. B¹, A. G¹, P. G¹, J. P¹, R. T¹, M. F¹, P. U¹, A. B², M. D², S², C. W², A. N.³
 P³, E. K³, K. E³, J. P. R⁴, N. L⁵, F. M⁶

Received 2003 October 23, accepted 2003 December 1, published 2004 January 23, ApJ 601:L163-L166, 2004 February 1

ABSTRACT

This Letter presents the first results of an observational campaign to study the Galactic center with *INTEGRAL*, the *International Gamma-Ray Astrophysics Laboratory*. Mosaicked images were constructed using data obtained with ISGRI, the soft gamma-ray instrument of the coded aperture IBIS imager, in the energy ranges 20–40 and 40–100 keV. These give a yet unseen view of the high-energy sources of this region in hard X-rays and gamma-rays with an angular resolution of 12' (FWHM). We report on the discovery of a source, IGR J1745.6–2901, coincident with the Galactic nucleus Sgr A* to within 0'.9. Located at R.A.(J2000.0) = 17^h45^m38^s.5, decl.(J2000.0) = –29°01'15'', the source is visible up to about 100 keV with a 20–100 keV luminosity at 8 kpc of $(2.89 \pm 0.41) \times 10^{35}$ ergs s^{–1}. Although the new *INTEGRAL* source cannot unequivocally be associated to the Galactic nucleus, this is the first report of significant hard X-ray emission from within the inner 10' of the Galaxy and a contribution from the Galactic supermassive black hole itself cannot be excluded.

Subject headings: black hole physics — Galaxy: center — Galaxy: nucleus — gamma-rays: observations — stars: neutron — X-rays: binaries

1. INTRODUCTION

From the prediction of the existence of a massive, compact source at the center of the Milky Way by Lynden-Bell & Rees (1971) to the discovery of such a source in the radio domain 3 years later by Balick & Brown (1974), and to the first detection of soft X-rays unmistakably attributable to it by *Chandra* in 1999 (Baganoff et al. 2003), the supermassive black hole candidate Sgr A* and the Galactic center (GC) region as a whole have been put under intense scrutiny for many years from radio wavelengths to gamma-rays. This has led to several discoveries, and advances in our understanding of the processes and interactions at the heart of the Milky Way. For example, it is now known that the soft X-ray emission (<10 keV) in the central 10' is heavily dominated by diffuse emission due primarily to hot gas (Koyama et al. 1996; Sidoli & Mereghetti 1999), and that only about 10% of the total emission in this energy range can be accounted for by X-ray point sources brighter than 10^{31} ergs s^{–1} (Muno et al. 2003a). Also, the contribution of point sources in this domain is about the same along the Galactic ridge as it is in the GC (Ebisawa et al. 2001).

The Galactic nuclear region consists of six primary components that give rise to an array of complex phenomena through their mutual interactions. These constituents are: Sgr A*, a supermassive black hole with a mass of around $3.5 \times 10^6 M_{\odot}$ (Schödel et al. 2002; Ghez et al. 2003a); the surrounding cluster of evolved and young stars; ionized gas streamers, some of which form a three armed spiral centered on Sgr A* known

as Sgr A West; a molecular dusty ring surrounding Sgr A West; diffuse hot gas; and a powerful supernova-like remnant known as Sgr A East (Melia & Falcke 2001). Furthermore, both the IR and the X-ray point source populations decreases in spatial density approximately as $1/R^2$, where R is the distance from the GC (Serabyn & Morris 1996; Muno et al. 2003a).

The first imaging observations of the GC in hard X-rays were performed by the X-ray telescope on Spacelab 2 in the range 2.5–20 keV with an angular resolution of 3'. A source located within 1'.1 of Sgr A* was detected (Skinner et al. 1987). Such a source was also detected by ART-P (8–20 keV) on the *Granat* satellite in 1990–1991 (Pavlin et al. 1994). At energies above 20 keV, only the coded mask instrument SIGMA/*Granat*, sensitive to energies above 35 keV, provided imaging capability with a ~20' angular resolution, allowing an exploration of the dense GC region. In spite of the deep 9×10^6 s SIGMA survey of the central parts of the galaxy performed between 1990 and 1997, only upper limits were set for the hard X- and gamma-ray emission from the neighborhood of Sgr A* at energies above 35 keV (Goldwurm et al. 1994; Goldoni et al. 1999). The derived low bolometric luminosity of the Galactic nucleus (GN), in contrast with the powerful output from active galactic nuclei or black hole binaries, has motivated the development of several models for radiatively inefficient accretion onto or ejection from Sgr A*. These models have been widely applied to other accreting systems (Melia & Falcke 2001).

A recent breakthrough discovery by *Chandra* and *XMM-Newton* is that Sgr A* is the source of powerful X-ray flares (Baganoff et al. 2001; Goldwurm et al. 2003a; Porquet et al. 2003) during which the soft X-ray luminosity can increase by factors of 50–180 over a period of up to 3 hr. Some of these X-ray flares feature a significant hardening of the spectrum up to photon indices of ~1. Furthermore, a very recent discovery with the Very Large Telescope NAOS/CONICA imager (Genzel et al. 2003) and the Keck telescope (Ghez et al. 2003b), is that Sgr A* is also the source of frequent IR flares. This activity could indicate the presence of an important population of non-thermal electrons in the vicinity of the black hole. These results have raised great interest in the possibility of observing hard X-rays from the GN, a measure of which may particularly shed light on the

*BASED ON OBSERVATIONS WITH INTEGRAL, AN ESA PROJECT WITH INSTRUMENTS AND SCIENCE DATA CENTRE FUNDED BY ESA MEMBER STATES (ESPECIALLY THE PI COUNTRIES: DENMARK, FRANCE, GERMANY, ITALY, SWITZERLAND, SPAIN), CZECH REPUBLIC AND POLAND, AND WITH THE PARTICIPATION OF RUSSIA AND THE USA.

¹ Service d'Astrophysique, DAPNIA/DSM/CEA, 91191 Gif-sur-Yvette, France; belanger@cea.fr

² IASF CNR, 00133 Roma, Italy; ubertini@rm.iasf.cnr.it

³ ESA, ESTEC, NL-2200 AG Noordwijk, The Netherlands; Christoph.Winkler@rssd.esa.int

⁴ ISDC, CH-1290 Versoix, Switzerland; ebisawa@obs.unige.ch

⁵ CESR, Toulouse Cedex 4, France; roques@Sigma-0.cesr.cnes.fr

⁶ DSRI, DK-2100 Copenhagen 0, Denmark; nl@dsri.dk

⁷ Physics Dept. and Stewart Observatory, University of Arizona, Tucson, AZ 85721, USA; melia@physics.arizona.edu

relative role of accretion and ejection in the Sgr A* system.

This Letter presents preliminary results of an observation campaign to study the GC at high energies with *INTEGRAL*, the *International Gamma-Ray Astrophysics Laboratory*, performed during the *INTEGRAL* Galactic Center Deep Exposure (GCDE) program in the spring of 2003. The focus is set on the inner $10'$ – $15'$ of the Galaxy and, more particularly, on the detection of a significant excess in this region. The results are based on data obtained with the *INTEGRAL* Soft Gamma-Ray Imager (ISGRI), the low-energy camera of the main Imager on Board the *INTEGRAL* Satellite (IBIS), proven to be accurate between 20 and 200 keV. The morphology of the central 2° in the 20–100 keV band and the derived positions of closeby high-energy sources are briefly discussed to elucidate the experimental context from which the results on the central source are drawn.

2. OBSERVATIONS AND RESULTS

INTEGRAL (Winkler 2003) is a European Space Agency observatory that began its mission on 2002 October 17 carrying four instruments. These consist of two main ones, IBIS (Ubertini et al. 2003) and SPI, the Spectrometer on *INTEGRAL* (Vedrenne et al. 2003); and two monitors, JEM-X (Lund et al. 2003) and OMC (Mas-Hesse et al. 2003).

The IBIS coded mask instrument is characterized by a wide field of view (FOV) of $29^\circ \times 29^\circ$, ($9^\circ \times 9^\circ$ fully coded), a point spread function (PSF) of $12'$ (FWHM), and a sensitivity over the energy range between 15 keV and 8 MeV. This sensitivity is achieved via two detector layers: ISGRI (Lebrun et al. 2003), an upper CdTe layer sensitive between 15 keV and 1 MeV with peak sensitivity between 15 and 200 keV, and PICsIT a bottom CsI layer, sensitive between 200 keV and 8 MeV.

We have analyzed IBIS/ISGRI data collected between 2003 February 28 and May 1 in a series of fixed pointings lasting about 37 minutes each. These include all GCDE data for which the pointings include the GN (~ 625 ks), and two Target of Opportunity observations (~ 475 ks). Data reduction was performed using the standard OSA 2 *INTEGRAL* Science Data Center analysis software (Goldwurm et al. 2003b), whose present version of the analysis procedures and calibration files do not allow for a full correction of systematic effects. Thus, in order not to over estimate the detection level of a source, taking into account fluctuations in the observation-dependent background noise levels as well as systematics, the significance was normalized to the fitted width of the distribution of individual significances in the image. This straight forward normalization procedure ensures that 68% of the distribution of significance values in the image are indeed contained within 1σ of the mean. Image reconstruction can be summarized as follows: from the events list for a pointing, subsets of events are selected according to energy bins. Each subset is used to build a detector image or shadowgram. Convolution of the shadowgram with a decoding array gives rise to a sky image containing the main peak of all sources in the FOV and their secondary lobes. Source identification and subtraction of secondary lobes results in the final reconstructed sky image. Fluxes are derived using *INTEGRAL* observations of the Crab Nebula performed just prior to the start of the GCDE, and luminosity calculations are based on a distance of 8 kpc to the GC assuming a Crab spectrum of a power law with photon index 2.12 and flux density at 1 keV of $8 \text{ ph cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$ (Bartlett 1994).

The maps of the GC shown in Figure 1 were constructed by summing the reconstructed images of 571 individual exposures. The total effective exposure time is about 8.5×10^5 s on the GN.




FIG. 1.— Milky Way's center seen in a $2^\circ \times 2^\circ$ field by the IBIS/ISGRI instrument in the energy ranges 20–40 keV (top) and 40–100 keV (bottom). Each image pixel size is equivalent to about 5 arcmin. Ten contour levels mark iso-significance linearly from about 4σ to 15σ .

of the Galaxy where ten contour levels mark iso-significance linearly from about 4σ up to 15σ , we can see what appear to be six distinct sources: 1E 1740.9–2942.7, KS 1741–293, A 1742–294, 1E 1743.1–2843, SLX 1744–299/300, whose nominal positions are marked by crosses and identification is still preliminary, and a source coincident with the radio position of Sgr A*. 1E 1740.7–2942 is a black hole candidate, KS 1741–293 and A 1742–294 are neutron star Low-Mass X-Ray Binary (LMXB) bursters, SLX 1744–299/300 are in fact two LMXBs separated by only $2.72'$, and 1E 1743.1–2843 is an X-ray source whose nature is still uncertain. In the 20–40 keV band, contours of the central source peak at the position of Sgr A* with a significance level of 8.7σ but are elongated to-

TABLE 1
INTEGRAL

Source name	Significance	Fitted position (R.A., decl.)	Offset ^a (arcmin)
1E 1740.7-2942	70.0	265.98, -29.74	0.32
KS 1741-293	18.9	266.23, -29.32	2.31
A 1742-294	15	266.52, -29.51	0.05
SLX 1744-299/300	9.6	266.86, -30.02	1.59
1E 1743.1-2843	9.2	266.59, -28.67	3.96
Sgr A*	8.7	266.41, -29.02	0.86

^aDistance between the fitted and nominal source positions

this transient neutron star LMXB burster system recently observed to have returned to an active state (Muno et al. 2003b) but could also be due to an uncorrected background structure. The central source is also marginally visible in the 40–100 keV band at a level of 4.7σ but without any contribution from the direction of GRS 1741.9–2853.

The position and flux of the central excess in the 20–40 keV map were determined by fitting the peaks with a function approximating the instrument’s PSF (Gros et al. 2003) in two different ways: (1) all the emission is attributed to one source and is fitted as such to determine its peak height and position, (2) the emission is attributed to two sources: a new source and GRS 1741.9–2853, whose position is then fixed to the one determined with the *Chandra* observatory (Muno et al. 2003b). Both of these involve a simultaneous fit of all the sources in the 2 deg² field of the GC. In the first case, we obtain a source position of R.A.(J2000.0)=17^h45^m22^s.5, decl.(J2000.0)=−28°58′17″, and a flux of about 5.4 mcrab or $(3.21 \pm 0.36) \times 10^{-11}$ ergs cm^{−2} s^{−1}. In the second, the position is 17^h45^m38^s.5, −29°01′15″, and the flux is about 3.2 mcrab or $(1.92 \pm 0.36) \times 10^{-11}$ ergs cm^{−2} s^{−1}. The central source’s 40–100 keV peak position is in very good agreement with the one determined using the second method outlined above, and since there is clearly no visible contribution from a neighboring source, the 40–100 keV flux was extracted at that position. The estimated error on the position is of about 4′ for a detection at the significance level of 8.7σ . These positions are respectively 4′6 and 0′9 from the radio position of Sgr A*. The results of the fine position determination for the six above mentioned GC sources are shown in Table 1. In Table 2, flux estimates for the central source are given adopting the result of the second method for position and flux determination. The hardness ratio (HR) — ratio of the count rate in the high-energy band over that in the low-energy band — for the detected excess is 0.90 ± 0.20 . As a possible indication of the nature of the detected excess, we can compare the values of the HR to the two brightest sources in the field. The BH candidate 1E 1740.7–2942 has a HR of 1.20 ± 0.03 , and the neutron star LMXB KS 1741–293 has a HR of 0.89 ± 0.08 .

The excess appears to be somewhat variable throughout the observations. In particular, we have detected a sudden increase in flux where the count rate rose by a factor of ~ 12 with respect to the mean for about 40 min. Figure 2 shows a portion of the light curve from the estimated 20–40 keV flux at the radio position of Sgr A*. Each data point corresponds to the average flux and statistical error from ~ 37 minute intervals during observations that lasted ~ 19 hr on 2003 April 6. Universal time is reported in *INTEGRAL* Julian days, i.e. days from 2000 January 1. The dashed line depicts the mean count rate over the




FIG. 2.— Portion of the total 20–40 keV lightcurve obtained by extracting the flux at the position of Sgr A* in the background corrected intensity map of each elementary exposure. The statistical errors are shown, and the dashed line corresponds to the mean count rate over the entire data set.

TABLE 2

Instrument	Energy (keV)	Energy Flux	
		(10^{-11} ergs cm ^{−2} s ^{−1})	(mCrab)
<i>INTEGRAL</i> (IGR 1745.6–2901)	20–40 ^a	1.92 ± 0.36	3.2 ± 0.6
	40–100 ^a	1.86 ± 0.40	3.4 ± 0.7
<i>XMM-Newton</i>	2–10 ^a	10.0	5.78
	20–40 ^b	0.61	1.03
	40–100 ^b	0.031	0.04
<i>BeppoSAX</i>	2–10 ^a	11.0	6.36
	20–40 ^c	0.89	1.49
	40–100 ^c	0.061	0.09

^aMeasured

^bExtrapolated using $kT = 7.2$ keV

^cExtrapolated using $kT = 8.0$ keV

significance of 5σ , bearing in mind that this significance may be hampered by systematic effects. A simultaneous increase in flux is seen with a lower significance in the 40–100 keV light curve. In what follows we discuss a number of possible explanations for this detected excess.

3. DISCUSSION AND CONCLUSION

In attempting to identify the excess at 17^h45^m38^s.5, −29°01′15″, we first consider which sources of high-energy emission can be excluded based solely on ISGRI’s error radius of 4′. There are four such sources: the radio arc located about 15′ from the GN; the non thermal X-ray filaments associated with it and the dense molecular cloud G0.13–0.13 (Yusef-Zadeh et al. 2002); GRS 1743–290, observed by SIGMA in 1991 (Goldwurm et al. 1994); and GRS 1741.9–2853 mentioned above and clearly unable to account for the bulk of the excess.

Second, we argue that this detection cannot be attributed to both diffuse emission and point sources as thus far observed at low X-ray energies within 8′–10′ of Sgr A*. This is so because it is incompatible with the extrapolation at higher energies of the flux integrated over this region as measured by *XMM-Newton* (A. Decourchelle 2003, private communication) and *BeppoSAX* (Sidoli et al. 1999). The analysis of 2001 *XMM-Newton* data in the 2–10 keV range consisted of integrating the total X-ray flux over a radius of 10′ around Sgr A*. The resulting spectrum is best fitted by a two-temperature plasma model with $kT_1 = 0.6$ keV for the soft component and

tions of the Sgr A complex over a radius of $8'$ around Sgr A* in the same energy band agree with *XMM-Newton*'s results. In this case the data are also best described by a two-temperature model with $kT_1=0.6$ keV and $kT_2=8$ keV. The extrapolation of the *XMM-Newton* and *BeppoSAX* results to higher energies was done using $kT=7.2$ and 8 keV, respectively. Both instruments' direct measurements of the diffuse X-ray flux are consistent with those of *Chandra* (Baganoff et al. 2003) and ART-P (Sunyaev et al. 1993). Table 2 reports the *XMM-Newton* and *BeppoSAX* measurements as well as their respective extrapolations. Note that the variability timescale of the *INTEGRAL* source equally excludes a major diffuse contribution and also strongly suggests that accelerated charged particles in the expanding shell of Sgr A East (Melia et al. 1998) cannot significantly contribute to the detected signal.

Third, since the detected excess cannot be fully accounted for by diffuse emission and is variable, it is reasonable to consider it a point source, IGR J1745.6–2901, of 20–100 keV luminosity $\sim 3 \times 10^{35}$ ergs s^{-1} . One known point source found within $4'$ of IGR J1745.6–2901 that could contribute to the signal is the eclipsing burster AX J1745.6–2901 (Maeda et al. 1996; Sakano et al. 2002). This source lies about $1'.3$ from Sgr A* and was detected in a high state only once, in 1994 (Sakano et al. 2002). The extrapolation of the flux measured in that state is marginally compatible with that of IGR J1745.6–2901, and this only in the low-energy range. The last confirmed detection of AX J1745.6–2901 was in 1997 by ASCA, when it was found in a much lower flux state. It may have recently been detected by *Chandra*. A timing analysis of the ASCA data on AX J1745.6–2901 revealed a modulation with a 40% decrease in flux and period of 8.356 ± 0.008 h (Maeda et al. 1996). 1997 ASCA data folded with that period show a slight dip, but since phase information could not be preserved over the 2.5 yr that separated the two observations, no unique solution for the period was found (Sakano et al. 2002). Folding the *INTEGRAL* data for IGR J1745.6–2901 with a period of 8.356 hr does not reveal any modulation.

Chandra observations of the GC performed on 2003 June 19 indicate that two sources, CXOGC J174540.0–290005 and CX-

OGC J174540.0–290014, found within $0'.4$ of Sgr A*, and another, CXOGC J174535.5–290124, lying $1'.3$ from Sgr A* and possibly the same as AX J1745.6–2901, had a combined intensity in the 2–8 keV band that was more than 30 times that of Sgr A* in its quiescent state on that day (F. K. Baganoff 2003, private communication). The GN was not particularly bright on June 19, but it must have been somewhat more active in May when an intense IR flare was observed to come from a region possibly very near the supermassive black hole's event horizon (Genzel et al. 2003).

It is noteworthy that the fairly hard X-ray source that ART-P detected in 1990 at the position of Sgr A*, had a 8–20 keV flux that varied between about 5 and 11 mcrab (Pavlinisky et al. 1994). This is consistent with our detection in the 20–40 keV energy range. We also recall the 2σ upper limits set by SIGMA on the 40–80 keV and 80–150 keV luminosity of the GN: 3.4×10^{35} and 2.8×10^{35} ergs s^{-1} respectively (Goldoni et al. 1999), both of which are compatible with our detection.

Finally, *INTEGRAL* has detected a very hard X-ray source at a position coincident to within $1'$ with Sgr A*. Because of IBIS's $12'$ angular resolution, the emission from IGR J1745.6–2901 cannot be attributed to one specific object and could be made up of several contributing sources found within a couple of arcminutes of Sgr A*. Nonetheless, this remains the first detection of emission at energies greater than 20 keV from the very close vicinity of the GN, and a contribution from Sgr A* itself cannot be excluded.

More *INTEGRAL* data are needed to better constrain the position, spectral shape, variability properties and the possibly multiple nature of IGR J1745.6–2901. We expect that the most constraining results will be provided by simultaneous observations in hard and soft X-rays with *INTEGRAL* and *XMM-Newton* or *Chandra*.

We gratefully thank François Lebrun for insightful suggestions and Anne Decourchelle for her analysis of *XMM-Newton* data on the GN.

REFERENCES

- Bartlett, L. M. 1994, Ph.D. thesis, Nasa Goddard Space Flight Center.
 Balick, B. & Brown, R. L. 1974, *ApJ*, 194, 265
 Baganoff, F. K. et al. 2001, *Nature*, 413, 45
 Baganoff, F. K. et al. 2003, *ApJ*, 591, 891
 Ebisawa, K. et al. 2001, *Science*, 293, 1633
 Genzel, R. et al. 2003, *Nature*, 425, 934
 Ghez, A. M. et al. 2003a, *ApJ*, accepted (astro-ph/0306130)
 Ghez, A. M. et al. 2003b, *ApJ*, in press (astro-ph/0309076)
 Goldoni, et al. 1999, *ApJ*, 38, 305
 Goldwurm, A. et al. 1994, *Nature*, 371, 589
 Goldwurm, A. et al. 2003a, *ApJ*, 584, 751
 Goldwurm, A. et al. 2003b, *A&A*, 411, L223
 Gros, A. et al. 2003, *A&A*, 411, L179
 Koyama, K. et al. *PASJ*, 48, 249
 Lebrun, F. et al. 2003, *A&A*, 411, L141
 Lund, N. et al. 2003, *A&A*, 411, L231
 Lynden-Bell, D. & Rees, M. J. 1971, *MNRAS*, 152, 461
 Maeda, Y. et al. 1996, *PASJ*, 48, 417
 Mas-Hesse, M. et al. 2003, *A&A*, 411, L461
 Melia, F. et al. 1998, *ApJ*, 508, 676
 Melia, F. & Falcke, H. 2001, *ARA&A*, 39, 309
 Muno, M. et al. 2003a, *ApJ*, 589, 225
 Muno, M. et al. 2003b, *ApJ*, in press (astro-ph/0307179)
 Pavlinsky, M. N., Grebenev, S. A. & Sunyaev, R. A. 1994, *ApJ*, 425, 110
 Porquet, D. et al. 2003, *A&A*, 407, L17
 Sakano, M. et al. 2002, *ApJS*, 138, 19
 Schödel, et al. 2002, *Nature*, 419, 694
 Serabyn, E. & Morris, M. 1996, *Nature*, 382, 602
 Sidoli, L. & Mereghetti, S. 1999, *A&A*, 349, L49
 Sidoli, L. et al. 1999, *ApJ*, 525, 215
 Skinner, G. K. et al. 1987, *Nature*, 330, 544
 Sunyaev, R. et al. 1993, *ApJ*, 407, 606
 Ubertini, P. et al. 2003, *A&A*, 411, L131
 Vedrenne, G. et al. 2003, *A&A*, 411, L63
 Winkler, C. 2003, *A&A*, 411, L1
 Yusef-Zadeh, F. et al. 2002, *ApJ*, 568, 121